

Molecular Diagnostics of Supernova Remnant Shocks

J. S. Lazendic¹, M. Wardle, A. J. Green

School of Physics, University of Sydney, NSW 2006, Australia

J. B. Whiteoak

Australia Telescope National Facility, CSIRO, PO Box 76, Epping NSW 1710, Australia

M. G. Burton

School of Physics, University of New South Wales, Sydney NSW 2052, Australia

Abstract.

We have undertaken a study of radio and infrared molecular-line emission towards several SNRs in order to investigate molecular signatures of SNR shocks, and to test models for OH(1720 MHz) maser production in SNRs. Here we present results on G349.7+0.2.

1. Introduction

Shocks driven by a supernova remnant (SNR) into a molecular cloud compress, accelerate and heat the gas, and can contribute to disruption of the molecular cloud. Despite the close association between SNRs and molecular clouds (Huang & Thaddeus 1986), establishing unambiguous cases of interaction has been difficult, mostly because of confusion resulting from unrelated clouds along the line of sight to the SNR and the possibility of chance alignment. Studies of SNR/molecular cloud interactions have been re-energized by the recent discovery of an association between OH(1720 MHz) maser emission and SNRs (see Koralesky et al. 1998, and references therein). This maser emission requires very specific conditions: gas density of $\sim 10^5 \text{ cm}^{-3}$, gas temperature of 50–125 K, and OH column density of $\sim 10^{16}$ – 10^{17} cm^{-2} (Lockett, Gauthier & Elitzur 1999). These conditions can be satisfied in the cooling gas behind a non-dissociative C-shock, irradiated by the X-ray flux from the SNR interior (Wardle 1999). The presence of OH(1720 MHz) masers in SNRs provides two crucial pieces of information for studying SNR shocks in molecular clouds: 1) the velocity of a cloud associated with an SNR, and 2) the precise location of the on-going interaction.

¹Current address: Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138




Figure 1. G349.7+0.2 is located at a distance of ~ 23 kpc, and is the third brightest galactic SNR at radio wavelengths. Five OH(1720 MHz) masers (marked with crosses) have been detected along the bright emission ridge of the SNR (Frail et al. 1996). The ^{12}CO 1–0 emission obtained with SEST is shown, integrated between 10 and 20 km s^{-1} (contours), and overlaid on the 18-cm radio continuum greyscale image obtained with ATCA with resolution of $9'' \times 5''$. The contour levels are: 16, 24, 32, 48, 63, 71, 78 K km s^{-1} . The dots mark the grid positions of the SEST observations.

2. Molecular gas towards the OH/SNR sites

For millimeter-line observations of molecular clouds interacting with SNRs, we used the Australia Telescope Mopra antenna and the Swedish-ESO Submillimetre Telescope (SEST). These observations had sufficient (sub-km/s) velocity resolution to associate the molecular gas with the OH masers, but suffered from poor spatial resolution ($20''$ – $50''$ for sources more than 8.5 kpc away). Observations of species such as ^{12}CO , ^{13}CO , CS, and HCO^+ were used to constrain the physical parameters of the ambient and shocked molecular gas. The observed molecular-line profiles imply the presence of a C-type shock in ambient gas with density of $\sim 10^4 \text{ cm}^{-3}$ and temperature of ~ 20 K. The shocked gas has a density of 10^5 – 10^6 cm^{-3} and temperature of $\gtrsim 40$ K. The latter properties of the molecular gas are consistent with the requirements for OH(1720 MHz) maser production.

3. H_2 emission

Millimeter-line observations were complemented by near-infrared (NIR) observations of H_2 emission with the University of New South Wales Infrared Fabry-Perot (UNSWIRF) narrow-band imaging spectrometer (Ryder et al. 1998), on the Anglo-Australian Telescope. UNSWIRF offers 75 km s^{-1} velocity resolution and $1''$ spatial resolution, which allows a determination of the relative positions of the OH masers and the shock-heated gas. We detected H_2 emission towards




Figure 2. (*Upper*) Shock-excited H₂ emission was detected towards the western edge of G349.2+0.7. The distribution of 2.12 μ m H₂ 1–0 S(1) emission (contours) is superimposed on the 18-cm ATCA image of G349.7+0.2. The contours are: 3.2, 9.7, 16.1 and 25.7 $\times 10^{-5}$ erg s⁻¹ cm⁻² sr⁻¹. H₂ observations revealed highly clumped molecular gas. An excellent correlation between the H₂ clumps and OH(1720 MHz) maser positions confirms the shock excitation of the masers. (*Lower*) OH absorption towards G349.2+0.7 was found at the radial velocity of the OH masers of $\sim +16$ km s⁻¹. The map of OH line-to-continuum ratio is shown, integrated between 10 and 20 km s⁻¹ (contours) overlaid on the 18-cm continuum image. The OH contours range from -0.01 to -0.03 Jy beam⁻¹ (beam is 15'' $\times$ 12'').

all four OH/SNR sources we observed (G359.1-0.5, G349.7+0.2, G357.7-0.1, G337.0-0.1). Shock excitation of H_2 is inferred either from the ratio of two H_2 2–1/1–0 S(1) lines, or from the correlation between H_2 and synchrotron radio emission. H_2 observations towards G349.7+0.2 are presented in Figure 2.

4. OH absorption

Theoretical models do not predict OH in any great abundance, because for temperatures above 400 K any OH formed from O and H_2 will be rapidly converted into H_2O . However, X-rays from the SNR interior can penetrate the surrounding molecular cloud, photo-ejecting electrons which collisionally excite H_2 . The subsequent radiative decay of H_2 contributes to a secondary UV radiation field, which converts part of H_2O back to OH after the temperature drops below 200 K (Wardle 1999).

The OH column necessary to produce OH(1720 MHz) masers can be observed in absorption at 18 cm against the background SNR radio continuum. We derived OH column density of $\sim 10^{16} \text{ cm}^{-2}$ towards three OH/SNR objects (W28, G349.2+0.7, G337.0-0.1) we observed using the Australia Telescope Compact Array (ATCA), which is consistent with the model for OH(1720 MHz) maser production. In Figure 2 we present results on G349.2+0.7. The maximum OH absorption occurs towards the OH masers and is well correlated with shocked H_2 emission. This is a clear indication that the enhancement of the OH gas is associated with shocked regions rather than background continuum intensity, as is usually found.

Acknowledgments. JSL was supported by the Australian Government International Postgraduate Research Scholarship. MW was supported by the Australian Research Council. The Australia Telescope is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. SEST is operated by the Swedish National Facility for Radio Astronomy, Onsala Space Observatory and by the European Southern Observatory (ESO).

References

- Frail, D. A., Goss, W. M., Reynoso, E. M., Giacani, E. B., Green, A. J. & Otrupcek, R. 1996, AJ, 111, 1651
- Huang, Y.-L. & Thaddeus, P. 1986, ApJ, 309, 804
- Koralesky, B., Frail, D. A., Goss, W. M., Claussen, M. J., & Green, A. J. 1998, AJ, 116, 1323
- Lazendic, J. S., Wardle, M., Burton, M. G., Yusef-Zadeh, F., Whiteoak, J. B., Green, A. J. & Ashley, M. C B. 2001, accepted for publication in MNRAS
- Lockett, P., Gauthier, E. & Elitzur, M. 1999, ApJ, 511, 235
- Ryder, S. D., Sun, Y.-S. Ashley, M. C B., Burton, M. G., Allen, L. E. & Storey, J. W. V. 1998, Publications of the Astronomical Society of Australia, 15, 228
- Wardle, M. 1999, ApJ, 525, L101